

Computer Science

Jerzy Świątek

Systems Modelling and Analysis

Choose yourself and new technologies


L.5.c Numerical optimization methods –
multidimensional search using derivatives



Project co-financed from the EU European Social Fund




Choice of the search direction



- Basis of search directions – non-gradient methods.
- Search directions based on gradient vectors – gradient-based methods.



Gradient based methods



$$x_{n+1} = x_n + \tau_n d_n$$
$$d_n = G(\nabla_x F(x_n))$$


τ_n – step size



The gradient descent method

$$x_{n+1} = x_n + \tau_n d_n$$


$$d_n = -\nabla_x F(x_n); \tau_n > 0, \lim_{n \rightarrow \infty} \tau_n = \tau, \sum_{n=0}^{\infty} \tau_n = \infty$$



$$\|x_{n+1} - x_n\| = \|\tau_n d_n\| < \varepsilon$$



The gradient descent method



$$x_{n+1} = x_n + \tau_n d_n$$


$d_n = -\nabla_x F(x_n)$, τ_n – optimal step size
along the direction d_n

$$\|x_{n+1} - x_n\| < \varepsilon$$



Newton's method

$$F(x) = \underbrace{F(x_0) + (x - x_0)^T \nabla_x F(x_0) + \frac{1}{2} (x - x_0)^T H(x_0) (x - x_0)}_{Q(x)} + O_3(\|x - x_0\|)$$



$$\nabla_x Q(x) = \nabla_x F(x_0) + H(x_0)(x^* - x_0) = O_S$$

$$x^* = x_0 - H^{-1}(x_0) \nabla_x F(x_0)$$

$$x_{n+1} = x_n - H^{-1}(x_n) \nabla_x F(x_n)$$



Fletcher-Reeves method of conjugate gradients

Step 0: $z_1 = x_0$, $s = 1$, $d_1 := -\nabla_x F(z_1)$

Step 1: $z_{s+1} := z_s + \tau_s d_s$

$\tau_s \rightarrow$ optimal step size along the direction d_s

If $\|\tau_s d_s\| < \varepsilon$ (STOP)

otherwise go to 2

Step 2: $d_{s+1} := -\nabla_x F(z_{s+1}) + \frac{\|\nabla_x F(z_{s+1})\|}{\|\nabla_x F(z_s)\|} d_s$

$s := s + 1$, go to 1

d_1, d_2, \dots, d_s – conjugate directions



Variable metric methods

$$z_1 = x_0$$

$$d_1 = -D_1 \nabla_x F(z_1) \quad D_1 = I$$

$$z_{s+1} = z_s + \tau_s d_s \quad \tau_s - \text{optimal step size along the direction } d_s$$

$$d_{s+1} = -D_{s+1} \nabla_x F(z_{s+1})$$

$$D_{s+1} = D_s + \frac{p_s p_s^T}{p_s^T q_s} - \frac{D_s q_s q_s^T D_s}{q_s^T D_s q_s},$$

$$p_s = \tau_s d_s$$

$$q_s = \nabla_x F(z_{s+1}) - \nabla_x F(z_s)$$

$$D_{s+1} \approx H^{-1}(x_{s+1})$$