

Computer Science

Jerzy Świątek

Systems Modelling and Analysis

Choose yourself and new technologies


L.17.b Numerical optimization methods –
multidimensional search without derivatives



Project co-financed from the EU European Social Fund





Choice of the search direction



- Basis of search directions – non-gradient methods.
- Search directions based on gradient vectors – gradient-based methods.



Method of Hooke and Jeeves with discrete steps



τ – step size
 $\alpha > 1$ exploratory step size
 $\beta \in (0,1)$ acceleration factor
 $\tau := \tau\beta$



Method of Hooke and Jeeves with discrete steps

Input data: $d_1, d_2, \dots, d_S, x_0, \tau, \varepsilon, \alpha, \beta$

Step 0: $z_1 := x_0, s = 1, n = 0$

Step 1: $z_{s+1} := z_s + \tau d_s$

 If $F(z_{s+1}) < F(z_s)$ then go to 2

 otherwise $z_{s+1} := z_s - \tau d_s$

 If $F(z_{s+1}) < F(z_s)$ then go to 2

 otherwise $z_{s+1} := z_s$

Step 2: If $s < S$, $s := s + 1$ then go to 1

 otherwise

 If $F(z_{S+1}) < F(z_1)$ then go to 3

$\tau := \tau\beta, x_{n+1} := x_n, z_s := x_n, n := n + 1, s = 1$ then go to 1


Step 3: If $\tau < \varepsilon$, STOP

 otherwise

$x_{n+1} := z_{S+1} + \alpha(z_{S+1} - z_1), n := n + 1, s := 1$ then go to 1



Method of Rosenbrock with discrete steps



τ – step size
 $\alpha > 1$ – exploratory step size acceleration
 $\beta \in (-1, 0)$ – acceleration factor
 $\tau_s := \tau_s \alpha$
 $\tau_s := \tau_s \beta$





Method of Rosenbrock with discrete steps

Input data: $d_1, d_2, \dots, d_S, x_0, \tau, \varepsilon, \alpha > 1, \beta \in (-1, 0)$

Step 0: $\tau_1 = \tau_2 = \dots = \tau_S = \tau, \delta_1 = \delta_2 = \dots = \delta_S = 0, z_1 = x_0, n = 0, s = 1$

Step 1: $z_{s+1} = z_s + \tau_s d_s, \delta_s = \delta_s + \tau_s$
 IF $F(z_{s+1}) < F(z_s)$ $\tau_s := \alpha \tau_s$ $s := s + 1$, THEN GO TO 2
 ELSE $F(z_{s+1}) \geq F(z_s)$ $\tau_s := \beta \tau_s$ $s := s + 1$ THEN GO TO 2

Step 2: IF $s < S$ $s := s + 1$ THEN GO TO 1

IF $F(z_{s+1}) < F(z_1)$ $z_1 := z_{s+1}$ $s := 1$ THEN GO TO 1

IF $F(z_{s+1}) = F(x_n)$

IF $n = 0$ change the initial solution


$x_{n+1} = z_{s+1}$

Step 3: ?

Step 4: Rotation of the basis of search directions



Method of Hooke and Jeeves using line searches





Method of Hooke and Jeeves using line searches

Input data: $d_1, d_2, \dots, d_S, x_0, \varepsilon$

Step 0: $z_s := x_0, n = 0, s = 1$

Step 1: $z_{s+1} := z_s + \tau_s d_s$

τ_s - optimal step size along the direction d_s

Step 2: If $s < S, s = s + 1$ then go to 1

If $\|z_{S+1} - z_1\| < \varepsilon$ - STOP

Step 3: $x_{n+1} := z_{S+1} + \tau d$


$\tau \rightarrow$ optimal step size along the direction d

$$d = \frac{z_{S+1} - z_1}{\|z_{S+1} - z_1\|} = \frac{\sum_{s=1}^{S+1} \tau_s d_s}{\|\sum_{s=1}^{S+1} \tau_s d_s\|}$$

$$n := n + 1, s = 1, z_1 = x_n$$



Method of Rosenbrock using line searches





Method of Rosenbrock using line searches

Input data: $d_1, d_2, \dots, d_S, x_0, \varepsilon$

Step 0: $z_1 := x_0, n = 0, s = 1$

Step 1: $z_{s+1} := z_s + \tau_s d_s$

τ_s - optimal step size along the direction d_s

Step 2: If $s < S, s := s + 1$ then go to 1

If $\|z_{s+1} - z_1\| < \varepsilon$ - STOP

Step 3: $a_s = \begin{cases} d_s & \tau_s = 0 \\ \sum_{j=s}^S \tau_j d_j & \tau_s \neq 0 \end{cases}$

$b_s = \begin{cases} a_s & s = 1 \\ a_s - \sum_{j=1}^{s-1} (a_j^T d'_j) d'_j \end{cases}$

$d'_s = \frac{b_s}{\|b_s\|} \quad s = 1, 2, \dots, S$


Step 4: $d_s := d'_s \quad s = 1, 2, \dots, S, n := n + 1, s = 1$ then go to 1



Powell's method – conjugate directions

d_1, d_2, \dots, d_s - conjugated directions,
 A – symmetric, positively defined matrix

$$d_i^T A d_j = \begin{cases} 0 & i \neq j \\ 1 & i = j \end{cases}$$






Powell's method – conjugate directions

$$F(x) = x^T Ax + b^T x + c$$

Optimizing along conjugate directions allows to reach solution after at most S steps.



Powell's method – conjugate directions




$x_1 = x_0 + \tau^* d$
 τ^* - optimal step size along the direction d from x_0
 $x_1' = x_0' + \tau^{*' } d'$
 $\tau^{*'}$ - optimal step size along the direction d' from x_0'

$d^T A d' = 0$
 d, d' - conjugated with respect A

$$d' = \frac{x_1' - x_1}{\|x_1' - x_1\|}$$



Powell's method





Powell's method

Input data: $d_1, d_2, \dots, d_S, x_0, \varepsilon$

Step 0: $z_1 := x_0 + \tau_S d_S, n := 0, \tau_S$ - optimal step size along the direction $d_S, s := 1$

Step 1: $z_{s+1} = z_s + \tau_s d_s$

τ_s - optimal step size along the direction d_s

Step 2: If $s < S, s := s + 1$ then go to 1

If $\|z_{S+1} - z_1\| < \varepsilon$ - STOP

Step 3: $x_{n+1} := z_{S+1}$

$$d := \frac{z_{S+1} - z_1}{\|z_{S+1} - z_1\|}$$

$z_1 := x_{n+1} + \tau d$ τ - optimal step size along the direction d

$d_s := d_{s+1}$ $s = 1, 2, \dots, S - 1$

$d_S := d$


$$\tau_{max} \rightarrow \max_{1 \leq s \leq S} \|z_{s+1} - z_s\| = \max_{1 \leq s \leq S} \tau_s$$

$$d_{max} = d \quad \Delta := \frac{\tau_m \Delta}{\|z_{S+1} - z_1\|} > 0.8$$



Nelder-Mead method

$x_1 x_2 \dots x_{S+1}$ - s -dimensional simplex



Initial simplex:

x_0, c

$d_j = [\quad]$

$$x_H \rightarrow F(x_H) = \max_{1 \leq s \leq S+1} F(x_s)$$

$$x_L \rightarrow F(x_L) = \min_{1 \leq s \leq S+1} F(x_s)$$

$$\bar{x} = \frac{1}{S} \sum_{s=1, s \neq H}^{S+1} x_s$$


$$a = \frac{c}{S\sqrt{2}} (\sqrt{S+1} + \sqrt{2} - 1)$$

$$b = \frac{c}{S\sqrt{2}} (\sqrt{S+1} - 1)$$

$$x_i = x_0 + d_j, x_{S+1} = x_0$$




Nelder-Mead method





Nelder-Mead method



Reflection

$$x^* = \bar{x} + \alpha(\bar{x} - x_H)$$

α – reflection coefficient

If $\alpha > 0$

$$F(x^*) < F(x_L)$$

Expansion

$$x^{**} = \bar{x} + \gamma(x^* - \bar{x}) \quad \gamma > 1$$

γ – contraction coefficient

If $F(x^*) > F(x_H)$

$$x^{***} = \bar{x} + \beta(x_H - \bar{x})$$

If $F(x^*) > \max_{\substack{1 \leq s \leq S+1 \\ s \neq H}} F(x_s)$

$$x^{***} = \bar{x} + \beta(x^* - \bar{x}) \quad \beta \in (0, 1)$$



Nelder-Mead method

Input data: x_0, c, ε

Step 0: x_1, x_2, \dots, x_{S+1} - initial simplex, $n = 0$

Step 1: $x_H \rightarrow F(x_H) = \max_{1 \leq s \leq S+1} F(x_s), x_L \rightarrow F(x_L) = \min_{1 \leq s \leq S+1} F(x_s)$

$$\bar{x} = \frac{1}{S} \sum_{\substack{s=1 \\ s \neq H}}^{S+1} x_s$$

Step 2: $x^* = \bar{x} + \alpha(\bar{x} - x_H)$

If $F(x^*) < F(x_L)$ $x^{**} = \bar{x} + \gamma(x^* - \bar{x})$ then go to 3
otherwise 4

Step 3: If $F(x^{**}) < F(x^*)$ $x_H = x^{**}, n = n + 1$ then go to 1
otherwise $x_H = x^*, n = n + 1$ then go to 1

Step 4: If $F(x^*) < \max_{\substack{1 \leq s \leq S+1 \\ s \neq H}} F(x_s)$ $x_H = x^*, n = n + 1$

Step 5: $x' - F(x') = \min\{F(x^*), F(x_H)\}$

$$x^{***} = \bar{x} + \beta(x' - \bar{x})$$

If $F(x^{***}) > F(x')$ $x_j = x_j + \frac{1}{2}(x_L - x_j), j = 1, 2, \dots, S + 1$ then go to 1

$x_H = x^{***}, n = n + 1$ then go to 1